

Review of α_s measurements at LEP 2

S. Martí i García

*Physics Department, University of Liverpool, Oxford St., Liverpool L69 7ZE, U.K.
e-mail: s.marti@cern.ch*

Abstract. Since 1995, LEP has steadily increased the center of mass energy of the colliding beams, from the M_Z resonance to 133, 161 and 172 GeV. New measurements of the strong coupling constant, α_s , at these energies have been performed by the LEP experiments, L3, ALEPH, OPAL and DELPHI. In this article, the new results are summarized, and combined with the previous LEP measurement of $\alpha_s(M_Z)$ in order to obtain an updated LEP average of $\alpha_s(M_Z) = 0.120 \pm 0.005$.

INTRODUCTION

Over the past two years (1995 and 1996), the RF system of LEP has been upgraded with the inclusion of new superconducting RF cavities. Thus the energy of the LEP e^- and e^+ beams has increased, giving data with center of mass energy of 133¹, 161 and 172 GeV. The analysis of the hadronic system produced in the $e^+e^- \rightarrow Z/\gamma^* \rightarrow q\bar{q}$ process allows the measurement of the running of the strong coupling constant, α_s , and the test of QCD predictions. Any deviation can be interpreted as a sign of new physics (e.g. production of light gluinos) appearing at the new energy domain accessible at LEP 2.

Theoretical predictions

It is well known that α_s is not a fixed but a running quantity. Its value depends on the physical energy scale (μ) of the process. The beta function describes the renormalization scale dependence of the strong coupling constant:

$$\mu \frac{d\alpha_s(\mu)}{d\mu} = -\frac{\beta_0}{2\pi} \alpha_s^2(\mu) - \frac{\beta_1}{4\pi^2} \alpha_s^3(\mu) + \mathcal{O}(\alpha_s^4) \quad (1)$$

¹⁾ The true energies were 130 and 136 GeV. However, as the integrated luminosity at these two energy points was the same, the data are usually combined and presented as data collected at the mean energy of 133 GeV.

Assuming SU(3) colour gauge symmetry: $\beta_0 = 11 - 2n_f/3$ and $\beta_1 = 51 - 19n_f/3$, with n_f being the number of active quark flavours. Solving equation 1, the α_s dependence with the scale goes as: $\ln(\mu^2/\Lambda^2)$, where Λ is the fundamental QCD scale.

Combination of individual results

It is a matter of discussion how the different measurements of α_s are combined, given that most of the theoretical uncertainties are correlated []. In this article, the average of a set of measurements, $x_i \pm \sigma_i$, is calculated as $\langle x \rangle = \Sigma x_i w_i$ (with weights w_i inversely proportional to squares of the error). The experimental error is obtained in a similar way, while the theoretical uncertainties are taken in average. The total error corresponds with the quadratic sum of the experimental and theoretical uncertainties.

MEASUREMENTS OF α_s AT LEP 1

The high statistics collected at the Z peak by the four LEP experiments (over 15 million of hadronic Z decays) has allowed $\alpha_s(M_Z)$ to be measured with high accuracy². All four LEP experiments have published α_s measurements with a wide range of methods. These can be summarized by: inclusive quantities as R_h and R_τ (the total hadronic decays of the Z and the τ), analysis of the global event shapes (thrust, broadness...), scaling violations [], three-jet rates, etc. Another set of important measurements was the test of the α_s universality for all the quark flavours []. Also important are the measurements using $q\bar{q}\gamma$ events, where the boost to the center of mass of the hadronic system allows measurements of the running of α_s [].

The global average of all LEP I measurements is: $\alpha_s(M_Z) = 0.121 \pm 0.005$, where the main contribution to the error is the theoretical uncertainty.

MEASUREMENTS OF α_s AT LEP 2

The α_s measurements at LEP 2 have new problems with respect to the measurements at LEP 1. The first is the limited statistics. So far, the integrated luminosity of the four experiments in LEP 2 is $\approx 100 \text{ pb}^{-1}$ (representing about 10,000 hadronic events). The second problem is the background, mainly from the radiative return to the Z events, and from $\gamma\gamma$ collisions. Another type of background comes from the W^+W^- pairs decaying into $q\bar{q}q\bar{q}$ and $q\bar{q}l\nu$, however this background is only present with LEP operating above the W^+W^-

²⁾ Due to this, most of the published results on α_s from other experiments at different energy scales are extrapolated to a common reference scale, usually taken as M_Z .

production threshold. Due to these problems, the total sample available for each individual experiment and energy point is about 400~1000 events.

Global event shapes

The LEP experiments obtained new results on $\alpha_s(\sqrt{s})$ by comparing the global event shape variables with the exact $\mathcal{O}(\alpha_s^2)$ QCD calculations plus either NLLA or resummed series. The new LEP 2 results for α_s are summarized in figure 1.




FIGURE 1. Measurements of α_s in LEP 2 at 133, 161 and 172 GeV. For each measurement, the errors quoted are the experimental and theoretical respectively. Dashed lines correspond to the QCD prediction assuming $\alpha_s(M_Z) = 0.118$.

The first measurement of $\alpha_s(133\text{GeV})$ was published by L3 [1]. In that analysis the distributions of thrust, scaled heavy jet mass and jet broadening variables were compared with resummed $\mathcal{O}(\alpha_s^2)$ QCD calculations. ALEPH [2] used the differential two jet rate with the Durham jet algorithm. In the DELPHI analysis [3] a similar set of event shape variables was used, plus the three jet rate with the Durham and the JADE algorithms. In the OPAL analysis [4] the distributions of $1 - T$, scaled heavy jet mass, jet broadening variables and the Durham differential two jet rate were fitted to the $\mathcal{O}(\alpha_s^2)$ +NLLA QCD predictions. The same event variables were used in the analysis of the 161

GeV data, providing the first $\alpha_s(161 \text{ GeV})$ LEP measurement [1].

The results at 172 GeV shown in figure 1 were provided by the QCD representatives of each experiment and are still preliminary.

The size of the nonperturbative effects in the event shape variables decreases when increasing the energy, as shown in a DELPHI study [2]. This is due to the convergence of the $\mathcal{O}(\alpha_s^2)$ perturbative calculations with the nonperturbative contribution described by a power series of α_s .

LEP average

In figure 2, the global LEP values for α_s are presented for each energy available at LEP. The lines corresponds to the running of α_s as obtained assuming $\alpha_s(M_Z) = 0.118 \pm 0.003$ and using the renormalization group equation. The new measurements are compatible with the expected values of α_s . Combining all values of $\alpha_s(M_Z)$ measured so far at LEP 1 and LEP 2, the new LEP average is:

$$\alpha_s(M_Z) = 0.120 \pm 0.005$$




FIGURE 2. Running of the strong coupling constant: measurements at different scales in LEP (M_Z , 133, 161 and 172 GeV by order). Only total errors are displayed.

COMPARISON WITH OTHER EXPERIMENTS

The LEP averages for α_s at 133, 161 and 172 GeV can be added to the plot showing α_s running (figure 3) over a wide range of scales [3]. With the new α_s results from LEP 2, the global average is found to be the same than in reference [3]:

$$\alpha_s(M_Z) = 0.118 \pm 0.003$$




FIGURE 3. Running of the strong coupling constant: measurements at different scales compared to the QCD prediction for $\alpha_s(M_Z) = 0.118 \pm 0.003$. The last three points correspond to the new LEP 2 measurements. Only total errors are displayed.

Acknowledgements

I would like to thank G. Cowan, D. Duchesneau, J. Fuster, L. del Pozo and D. Wicke for their help in preparing the talk. Also to J.M. Hernández, M. Martínez, J. Terrón and J. Roldán for the pleasant week in Chicago. I apologise to everybody whose contribution in this vast field I failed to mention.

REFERENCES

1. M. Schmelling XXVIII ICHEP, Warsaw July 1996, hep-ex/9701002
2. ALEPH coll. Phys. Lett. B357 (1995) 487
3. DELPHI coll. CERN-PPE/96-185
4. DELPHI coll. Phys. Lett. B307 (1993) 221
5. DELPHI coll. Zeit. Phys. C70 (1996) 179
6. ALEPH coll. CERN-PPE/96-186
7. S. Martí i García, hep-ex/9511004
8. L3 coll. Contribution to ICHEP 96, Warsaw, pa04-021
9. L3 coll. Phys. Lett. B371 (1996) 137
10. ALEPH coll. Zeit. Phys. C73 (1997) 409
11. DELPHI coll. Z. Phys. C73 (1997) 229
12. OPAL coll. Z. Phys. C72 (1996) 191
13. OPAL coll. CERN-PPE/97-015
14. K. Hamacher. Proceedings of QCD 96 conference, Montpellier.